

## Other Criteria for Queries in IDdx

### **Disease Categories**

Arthropod-Borne  
Cestodes  
Childhood Infections  
Foodborne Intoxications  
Gastroenteritis--Inflammatory  
Gastroenteritis--Noninflammatory  
Immune-Related  
Intestinal Nematodes  
Localized Infections  
Person-To-Person  
Sapronoses  
Sexually-Transmitted  
Tissue Nematodes  
Trematodes  
Zoonoses  
Exclude Localized Infections

### **Epidemiological Factors**

**Entry:** Animal Bite; Ingest; Inhale; Needle; Sex; Skin/Eye; Swim

**Source:** Dairy; Edible Plant; Excreta; Fecal-Oral; Food; Meat; Patient; Seafood-Shellfish; Soil; Tissue; Water

**Vector:** Biting Flies; Fleas; Lice; Mites; Mosquitoes; Ticks

**Reservoir:** Birds; Cats; Cattle; Deer; Dogs; Fish; Horses; Human; Monkeys; Rabbits; Rodents; Swine; Wildlife

### **Other Criteria**

**Endemic:** Not in North America; Australia and South Pacific; East Asia; India and South Asia; Southeast Asia; North America; Mexico, Central America and Caribbean; Temperate South America; Tropical South America; North Africa; Southern Africa; West Africa; Central Africa; East Africa; Middle East; Western Europe; Eastern Europe and Former USSR

**Incubation Period:** 0-6 hours; 7-24 hours; 24-48 hours; 2-14 days; 2-8 weeks; 2-24 months; >2 years

**Acuity:** Acute-Severe; Acute-Moderate; Subacute/Chronic

**Immunocompromised at Risk**

**Bioterrorism Diseases**